

Citation for published version:

Dunhill, A & Wills, M 2015, 'Geographic range did not confer resilience to extinction in terrestrial vertebrates at the end-Triassic crisis', *Nature Communications*, vol. 6, 7980, pp. 1-8. <https://doi.org/10.1038/ncomms8980>

DOI:

[10.1038/ncomms8980](https://doi.org/10.1038/ncomms8980)

Publication date:

2015

Document Version

Peer reviewed version

[Link to publication](#)

Publisher Rights

Unspecified

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Geographic range did not confer resilience to extinction in terrestrial vertebrates at the end-Triassic crisis

Alexander M. Dunhill^{1,2} and Matthew A. Wills¹

¹Milner Centre for Evolution, University of Bath, Claverton Down, Bath, BA2 7AY, UK

²School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, UK,
a.dunhill@leeds.ac.uk

Rates of extinction vary greatly through geological time, with losses particularly concentrated in mass extinctions. Species duration at other times varies greatly, but the reasons for this are unclear. Geographical range correlates with lineage duration amongst marine invertebrates, but it is less clear how far this generality extends to other groups in other habitats. It is also unclear whether a wide geographical distribution makes groups more likely to survive mass extinctions. Here, we test for extinction selectivity amongst terrestrial vertebrates across the end-Triassic event. We demonstrate that terrestrial vertebrate clades with larger geographical ranges were more resilient to extinction than those with smaller ranges throughout the Triassic and Jurassic. However, this relationship weakened with increasing proximity to the end-Triassic mass extinction, breaking down altogether across the event itself. We demonstrate that these findings are not a function of sampling biases; a perennial issue in studies of this kind.

Is it possible to make generalisations about which clades and higher taxa are most likely to go extinct on geological timescales? Geographic range is often claimed to be a determinant of extinction vulnerability¹⁻⁶. Groups with restricted ranges are widely believed to suffer extinction more frequently than those with broader ranges¹ because the latter are thought to

26 be afforded some resilience to regional environmental perturbations^{7,8}. During global biotic
27 crises, by contrast, there is reasoned to be no such selectivity^{2,6} since widespread
28 environmental disturbances simultaneously affect both wide and narrow ranging taxa on
29 global scales^{1,3,6}. Surprisingly, the effect of geographic range on extinction risk has not been
30 tested comprehensively for the terrestrial fossil record, with a striking paucity of studies on
31 vertebrates of any kind. Most published studies focus on marine invertebrates^{3-6,8-10}, and
32 despite geographic range being used as a major criterion for assessing the extinction risk of
33 modern terrestrial species¹¹, it is unclear that the findings from these fossil taxa can be
34 extended to all groups in all major habitats. Moreover, little is known about the difference
35 between intervals with background levels of extinction and those characterised as mass
36 extinctions³. The only way to answer such questions is to utilise fossil evidence of past
37 organismal distributions^{5,12}.

38 The Triassic to Jurassic is a critically important period in terrestrial vertebrate
39 evolution^{13,14}. In particular, it follows the largest of all mass extinctions, the Permo-Triassic
40 (P/T) ‘great dying’¹³⁻¹⁵. Many terrestrial vertebrate lineages originated in the aftermath of the
41 P/T event, but were subsequently subjected to major changes in terrestrial ecosystems
42 throughout the ensuing Triassic and Jurassic. These changes included the gradual aridification
43 of Pangaea¹⁶, as well as its initial rifting and fragmentation¹⁷ allied to the eruption of the
44 Central Atlantic Magmatic Province (CAMP)^{18,19}. This culminated in the end-Triassic mass
45 extinction event^{16,20}, which saw the demise of numerous amphibian and reptile groups prior
46 to the subsequent rise to dominance of the dinosaurs^{13,16,21-23}.

47 The significant vertebrate faunal turnover throughout the Triassic and Jurassic (lying
48 either side of the end Triassic mass extinction) make this an ideal period in which to study
49 extinction selectivity. We therefore collated palaeobiogeographical and stratigraphic
50 distributional data²⁴ for Triassic and Jurassic terrestrial vertebrate clades in order to ask three

51 questions. 1. Is there is a relationship between palaeobiogeographical distribution and the risk
52 of extinction during ‘normal’ times? 2. Does any such relationship disappear across the end-
53 Triassic mass extinction? 3. Can any of these findings be attributed to sampling biases?

54 We find that wider geographical range conferred greater resilience to extinction in
55 terrestrial vertebrate groups throughout most of the Triassic and Jurassic. However, this
56 insurance weakened towards the end-Triassic mass extinction, and was imperceptible across
57 the event itself. Major sampling biases were discounted as the cause of these patterns.

58

59 **Results and Discussion**

60 **Geographic range and diversification rates are correlated.** Diversification rates and
61 changes in geographic range at the clade level are positively and strongly correlated across all
62 time bins (Figs. 1 and 2 and Supplementary Table 1), with weaker (but still mostly
63 significant) correlations when the data are subdivided into Epochs (Fig. 2 and Supplementary
64 Table 1). The strong correlation between changes in geographic range and diversification rate
65 across all time intervals (and at Epoch level and within stage-level time bins) confirms that
66 increasing range size coincides with increasing diversity, whilst range size reductions
67 typically attend diversity reductions. Taxa with larger geographic ranges are therefore more
68 likely to exhibit increasing diversity and lower extinction rates than those with smaller
69 ranges. Greater rates of origination might also be expected to result from more extensive
70 ranges; firstly because large ranges are more likely to be fragmented into peripheral isolates,
71 secondly because larger ranges encompass a greater diversity of environments and selective
72 forces^{25,26}. This pattern is the opposite of that proposed for the marine invertebrate fauna,
73 where taxa with narrow ranges show higher origination rates^{6,27}. The results from our
74 geographic range data sets, both raw convex hull and standardized mean great circle distances

75 (Fig. 1), are similar, demonstrating that our findings are not contingent upon the precise
76 methodology used.

77

78 **This relationship breaks down across the mass extinction.** Strikingly, rates of geographic
79 range change and diversification are not significantly correlated immediately before the end-
80 Triassic extinction (during the Rhaetian), whereas this correlation is significant even in the
81 Carnian and Norian stages that immediately precede it (Fig. 2 and Supplementary Table 1).
82 Hence, diversification rate becomes decoupled from range size change rate across the
83 extinction boundary and the insurance against extinction offered by larger geographic ranges
84 at other times disappears. During the end-Triassic mass extinction event, relatively
85 widespread groups are as likely to suffer high levels of extinction as groups with narrower
86 geographic ranges. For example, Phytosauria and Theropoda have similar, geographic ranges
87 in the Rhaetian (Fig. 3). However, phytosaurs suffer complete extinction at the end-Triassic,
88 whereas theropod diversity remains stable across the boundary and into the Hettangian, even
89 whilst undergoing significant range expansion (Fig. 3). Of the time intervals that do not show
90 significant correlation between diversification rate and geographic range change rate, all have
91 very small sample sizes apart from the Rhaetian. The standardised mean great circle distance
92 (GCD) range metrics show less correlation with diversification rate than the raw convex hull
93 metric (Supplementary Table 1). This is unsurprising since a reduced sample size leads to a
94 reduction in statistical power. However, many of these non-significant correlations still have
95 relatively high correlation coefficients and are approaching significance, whereas the
96 Rhaetian results are clearly different (with negative coefficients that are far from significant)
97 (Supplementary Table 1). Our temporal divisions are very much longer (2.0 to 18.9
98 Myrs)^{28,29} than the extinction event²⁰, which occurred in pulses over a period of less than 1.0
99 Myr¹⁹. Effects are therefore time averaged, meaning that the breakdown of the relationship

between diversification rate and the rate of geographic range change in the very end Triassic (an interval known to contain a major biotic crisis) is even more striking.

Sampling biases have little effect on our results. We observed several significant bivariate correlations between diversification rate, geographic range change rate and various putative sampling proxies detrended using first differences (Supplementary Tables 2-3). However, multiple regression models identified the changes in geographic range rate as the dominant variable influencing diversification rate (Table 1 and Supplementary Tables 4-9).

Although certain parts of the Triassic-Jurassic are reputed to have a poor terrestrial fossil record (i.e. Ladinian, Toarcian-Mid Jurassic)^{13,20,30-32}, the Late Triassic possesses one of the best^{23,32} (Fig. 4). We see positive correlations between land area and geographic range and a negative correlation between sea level and range in the GCD data, but also positive correlations between sea level and geographic range and between sea level and diversification rate in the convex hull data. This suggests that greater land area and lower sea levels may result in greater geographic ranges amongst terrestrial organisms. However, the negative correlation between standardised range and sea level (and the lack of correlation between diversification rate and land area) suggests that while expanding landmasses might be expected to induce the expansion of terrestrial ranges and increase diversification, climatic and other effects complicate this relationship. Indeed, the typically harsh environments in deep continental interiors may constrain many ranges. Specifically, lethally hot temperatures, particularly in the Early Triassic, may have limited or even excluded diversity in equatorial regions³³. Individual lineage ranges correlate with total ranges across time bins, and both appear to reflect the same underlying pattern. The considerable variation between individual lineage ranges within each bin (coupled with the fact that the standardised range metrics still show some correlation with total range, albeit weaker than the raw range metrics) indicates

that range estimates are not governed straightforwardly by sampling intensity (Figs. 1 and 3). Moreover, the absence of terrestrial vertebrates from equatorial regions is entirely plausible during the climatic greenhouse of the Early Triassic³³. Groups with broader geographical distributions are likely to be subject to a wider range of selective pressures and the peripheral isolation of sub groups; both factors favouring speciation and increasing diversity^{25,26}. However, the weak correlation observed between changes in fossil abundance and both diversification rate and geographic range change may be representative of sampling bias. It is also possible that a greater investment of research effort in more abundant fossil groups may have resulted in increased taxonomic splitting³⁴.

We also observed significant pairwise correlations between both raw and standardised geographic range change and diversification rate on one hand, and changes in numbers of formations on the other. Although formation counts are regarded as effective sampling proxies for terrestrial data sets³⁵, redundancy between sampling proxies and diversity metrics (arising from the probable non-independence of formation and fossil content) remains problematic³⁶⁻³⁸. In practice, the level of this redundancy is likely to be minimised by the use of all terrestrial vertebrate-bearing formations^{39,40}, rather than by adopting a stricter count of only those formations containing a particular group of terrestrial vertebrate fossils^{31,41-43}. However, standardisation of geographic range data results in the removal of significant correlations between range size and fossil abundance, coupled with a weakening of the correlation between range size and total range size (i.e. standardising geographic range calculations to a constant sample size across all lineages in each time bin appears to remove putative sampling effects). By contrast, standardising range data does not affect the correlation between range size and formation counts. This last relationship may therefore arise from redundancy³⁶, rather than being a temporal bias resulting from variation

in the amount of preserved sedimentary rock (and concomitant intensity of sampling) through geological time.

The multiple regression models show that geographic range change is the dominant driver of diversification rate through the Triassic-Jurassic, to the exclusion of all the sampling proxies in the model using convex hulls, but with total range and sea level showing some influence in the standardised mean great circle distance models. This suggests that, although sampling biases are a perennial issue in fossil data sets, the link between changes in geographic range and diversification rate appear robust, despite the patchy nature of the vertebrate fossil record.

Implications for extinction studies. We demonstrate that broad geographic range conferred insurance against extinction on major clades of terrestrial vertebrates during periods with background levels of extinction. In line with marine invertebrate studies across the same biotic crisis⁶ and at other times in the Phanerozoic³, this insurance disappeared during the end-Triassic mass extinction. However, these results are in marked contrast to patterns reported for marine invertebrates at the Cretaceous-Paleogene extinction^{9,44} (where groups with larger geographic ranges retain increased resilience to extinction across the crisis than those with smaller ranges). It is reasonably common to observe discrete macroevolutionary patterns in different higher taxa or across different major habitats. Notable examples include the incongruence between terrestrial and marine Phanerozoic diversity curves⁴⁵⁻⁴⁷, variations in the apparent force of Cope's rule *sensu lato* in different higher taxa⁴⁸⁻⁵⁰, and variations in the relationships between body size, population density and fecundity across clades^{51,52}. However, the differences between patterns observed at major extinction events may result from differences between the particular driving forces of the crises themselves. These differences highlight the need for greater numbers of large-scale, comparative studies before

attempting to make macroevolutionary generalisations. Fortunately, the expansion and refinement of resources such as the Paleobiology Database, coupled with quantitative tools such as GIS, have great potential for testing temporal and spatial macroevolutionary patterns. Equally, palaeontological data provides a broader perspective on the current biodiversity crisis. Specifically, it enables deep time tests of the purported relationships between present day extinction susceptibility and geographical range size^{53,54}, latitudinal distribution⁵⁵, niche breadth⁵⁶ and body size^{57,58}.

Methods

Fossil occurrence data. Stage level occurrence data for Triassic-Jurassic terrestrial vertebrates were initially downloaded from the Paleobiology Database²⁴ (<https://paleobiodb.org>) on 4th February, 2013 (last accessed 20th April, 2015) and, after pre-processing, consisted of 3507 occurrences of 857 genera (see Supplementary Note 1 for Paleobiology Database download specifications). Terrestrial vertebrate occurrences from marine deposits were not included as they would not give a true representation of geographic range. Ichnogenera and other form taxa were then removed from the data set as they could not be assigned accurately to parent genera. Marine tetrapod taxa recorded in terrestrial formations were also removed. Generic indeterminate taxa and taxa with uncertain generic assignments (i.e. *aff.*, *cf.*, *ex gr.*, *sensu lato*, ?) were also excluded. Although these exclusions inevitably resulted in an underestimation of the geographic range of higher taxonomic groups, it would be inappropriate to compare ranges constructed from taxa of uncertain affiliation with rates of generic extinction, origination, and diversification, which cannot include these indeterminate occurrences.

Fossil occurrences were vetted for synonymy and outdated taxonomy and sorted into higher taxonomic groups according to phylogenetic and ecological relationships within the

constraints of reasonable sample sizes (see Supplementary Note 1 for detailed classification of taxa). As with all higher taxonomic classification, groups were not directly comparable. This is an unavoidable problem unless working at the species or, to a lesser extent, the generic level. Two datasets were compiled: data₁ and data₂ (Supplementary Fig. 1). Data₁ (Supplementary Data 1) contained a number of paraphyletic assemblages where basal taxa of particular lineages were grouped together to form a paraphyletic ‘stem’ assemblage (e.g. Archosauriformes, basal Cynodontia, Dinosauromorpha) (Supplementary Fig. 1). Since the inclusion of paraphyletic groups is arguably problematic (they do not represent ‘true’ evolutionary groups), a second dataset excluding all parapylla was also prepared. In data₂, the paraphyletic assemblages were collapsed into smaller, monophyletic family-level groups wherever possible (Supplementary Fig. 1). The two data sets correlated very closely in terms of both geographic range change rate and diversification rate. All analyses in the manuscript therefore refer exclusively to data₁.

Fossil occurrences were binned at the stratigraphic stage level. Any occurrences not assigned to a stage were attached to the stage, or range of stages, corresponding to the formation from which they were recovered. Fossil occurrences that were assigned to more than one stage were randomly assigned to a single stage within their given range, a process that was repeated 1,000 times to obtain a mean value. This method avoided either the artificial inflation or deflation of taxonomic richness in any given stage compared to maximum or minimum diversity values.

Sampling and environmental proxy data. *Non-marine area*⁵⁹: A mean estimate of the continental landmass for each Stage. It was anticipated that geographic range would correlate positively with land area as the area of terrestrial habitat creates an upper limit for the geographic range of terrestrial organisms. These measurements were derived from an

independent source⁵⁹, and were subject to different definitions of Stage-level boundaries than the fossil occurrence data set, which used the Geological Time Scale 2012⁶⁰.

*Sea level*⁶¹: A mean estimate of relative sea level for each Stage. It was expected that geographic range would correlate inversely with mean sea level, as higher sea level would result in less terrestrial landmass for terrestrial organisms to inhabit. As with the non-marine area measurements, the sea level averages were obtained from an independent source⁶¹ and are subject to different definitions of Stage-level boundaries from the fossil occurrence data set⁶⁰.

*Terrestrial formations*²⁴: Formation counts are widely regarded as effective sampling proxies for the terrestrial fossil record^{31,35,42,62}. It is still unclear if this is true, as formation counts probably share a common signal with fossil occurrence data (i.e. formations are not independent from their fossil content^{36,37}). However, given the lack of comprehensive data on global sedimentary rock outcrop area, formation counts offer the only possible metric of global rock availability. In this analysis, redundancy was minimised by using a total count of terrestrial tetrapod bearing formations, rather than a strict count of group-specific bearing formations. There is also an argument for redundancy between formation counts and geographic range, as a taxon that is genuinely wide ranging is more likely to be present in more formations across the globe than a taxon with a small geographic range. Such possibilities were explored using multiple regression models.

*Fossil abundance*²⁴: Fossil abundance per time period serves as a proxy for human sampling and collecting effort. However, there is danger of circularity, as palaeontologists will be more likely to collect from formations yielding a higher number of fossils^{38,63}. Therefore, fossil abundance may be more representative of preservational factors or true biological abundance, rather than a measure of human sampling effort.

Total geographic range: It is reasonable to assume that vertebrates were not genuinely absent from large areas of the globe through parts of the Triassic-Jurassic. Therefore, if total geographic range (i.e. the total geographic range of all tetrapod occurrences per time bin) were to correlate strongly with the geographic ranges of individual fossil groups, it would indicate that the pattern of geographic range through the study period is controlled by spatial sampling rather than reflecting a biological pattern.

Analysis. Palaeogeographic ranges were constructed by converting modern fossil occurrence coordinates to palaeocoordinates using *PointTracker*⁶⁴. Palaeogeographic ranges were constructed using two methods: (i) in *ArcGIS v10.1* using convex hulls around the palaeogeographic occurrences for each group^{65,66}; and (ii) using mean great circle distances (GCD) between palaeogeographic occurrences with sample sizes standardised to 5 and 10 occurrences per lineage per time bin and replicated 1,000 times to obtain a mean value. GCD distances were calculated using the spherical law of cosines (as an acceptable approximation of the Haversine formula for terrestrial calculations).

Per lineage origination (O_r) and extinction (E_r) rates were calculated using the methodology of Foote⁶⁷ and modified by Foote⁶⁸:

$$(1) \quad O_r = -\ln \frac{N_{bt}}{N_{ft} + N_{bt}}$$

$$(2) \quad E_r = -\ln \frac{N_{bt}}{N_{bL} + N_{bt}}$$

Where N_{bt} = number of range-through taxa, N_{ft} = number of taxa that originate within time bin and cross top boundary of time bin, N_{bL} = number of taxa that cross bottom boundary of time bin but make their last appearance in time bin. Rates were not expressed relative to time bin duration; although this may cause underestimation of rates in shorter time bins relative to

longer time bins, Foote⁶⁹ demonstrated that both extinction and origination are pulsed rather than spread throughout time intervals. All analyses were carried out at the generic level.

No significant correlations were detected between geographic range change and extinction rate or between geographic range change and origination rate (Supplementary Table 10). The absence of significant correlations between origination/extinction rates and change in geographic range could be regarded as somewhat surprising, but this result is a function of the paucity of data for the rate calculations. However, the extinction and origination rate calculations rely on taxa that range-through three consecutive time bins⁶⁷, which are scarce in this data set because of the patchiness of the terrestrial fossil record and the long durations of the time bins. Therefore, it was judged that a metric of diversification calculated from generic range data offered a more robust picture of biotic change. Diversification rate (D_r) and geographic change rate (R_r) were calculated using a metric modified from Rode and Lieberman⁷⁰:

$$(3) \quad D_r = (\ln D_1 - \ln D_0)$$

$$(4) \quad R_r = (\ln R_1 - \ln R_0)$$

where D_1 = ranged-through diversity calculated from first and last appearances for current time interval, D_0 = ranged-through diversity calculated from first and last appearances for the previous time interval, R_1 = geographic range in time interval, R_0 = geographic range in previous time interval.

Relationships between geographic range change and generic diversification rates within clades were tested using pairwise Spearman's rank-order correlation tests. Putative sampling biases were investigated using both pairwise correlation and multiple regression models, with diversification rate as the dependent variable and geographic range and various sampling proxies as independent variables. First differencing was used to detrend the

sampling proxy data prior to comparison with the biotic rate data. All statistical analyses were carried out in R 3.1.1.

References

- 1 Barnosky, A. D. *et al.* Has the Earth's sixth mass extinction already arrived? *Nature* **471**, 51-57, doi:10.1038/nature09678 (2011).
- 2 Jablonski, D. Mass extinctions and macroevolution. *Paleobiology* **31**, 192-210 (2005).
- 3 Payne, J. L. & Finnegan, S. The effect of geographic range on extinction risk during background and mass extinction. *Proceedings of the National Academy of Sciences of the United States of America* **104**, 10506-10511, doi:10.1073/pnas.0701257104 (2007).
- 4 Powell, M. G. Geographic range and genus longevity of late Paleozoic brachiopods. *Paleobiology* **33**, 530-546, doi:10.1666/07011.1 (2007).
- 5 Rode, A. L. & Lieberman, B. S. Using GIS to unlock the interactions between biogeography, environment, and evolution in Middle and Late Devonian brachiopods and bivalves. *Palaeogeography, Palaeoclimatology, Palaeoecology* **211**, 345-359, doi:10.1016/j.palaeo.2004.05.013 (2004).
- 6 Kiessling, W. & Aberhan, M. Geographical distribution and extinction risk: lessons from Triassic-Jurassic marine benthic organisms. *Journal of Biogeography* **34**, 1473-1489, doi:10.1111/j.1365-2699.2007.01709.x (2007).
- 7 Foote, M., Crampton, J. S., Beu, A. G. & Cooper, R. A. On the bidirectional relationship between geographic range and taxonomic duration. *Paleobiology* **34**, 421-433, doi:10.1666/08023.1 (2008).

- 318 8 Nürnberg, S. & Aberhan, M. Habitat breadth and geographic range predict diversity
319 dynamics in marine Mesozoic bivalves. *Paleobiology* **39**, 360-372, doi:10.1666/12047
320 (2013).
- 321 9 Jablonski, D. Extinction and the spatial dynamics of biodiversity. *Proceedings of the*
322 *National Academy of Sciences of the United States of America* **105**, 11528-11535,
323 doi:10.1073/pnas.0801919105 (2008).
- 324 10 Jablonski, D. & Roy, K. Geographical range and speciation in fossil and living
325 molluscs. *Proceedings of the Royal Society B* **270**, 401-406 (2003).
- 326 11 Mace, G. M. *et al.* Quantification of Extinction Risk: IUCN's System for Classifying
327 Threatened SpeciesCuantificación del Riesgo de Extinción: Sistema de la UICN para
328 la Clasificación de Especies Amenazadas. *Conservation Biology* **22**, 1424-1442,
329 doi:10.1111/j.1523-1739.2008.01044.x (2008).
- 330 12 Valentine, J. W., Jablonski, D., Krug, A. Z. & Berke, S. K. The sampling and
331 estimation of marine paleodiversity patterns: implications of a Pliocene model.
332 *Paleobiology* **39**, 1-20, doi:10.1666/0094-8373-39.1.1 (2013).
- 333 13 Brusatte, S. L., Benton, M. J., Desojo, J. B. & Langer, M. C. The higher-level
334 phylogeny of Archosauria (Tetrapoda: Diapsida). *Journal of Systematic*
335 *Palaeontology* **8**, 3-47, doi:10.1080/14772010903537732 (2010).
- 336 14 Ezcurra, M. D. Biogeography of Triassic tetrapods: evidence for provincialism and
337 driven sympatric cladogenesis in the early evolution of modern tetrapod lineages.
338 *Proceedings of the Royal Society B: Biological Sciences* **277**, 2547-2552,
339 doi:10.1098/rspb.2010.0508 (2010).

- 340 15 Ruta, M., Botha-Brink, J., Mitchell, S. A. & Benton, M. J. The radiation of cynodonts
341 and the ground plan of mammalian morphological diversity. *Proceedings of the Royal*
342 *Society B: Biological Sciences* **280**, doi:10.1098/rspb.2013.1865 (2013).
- 343 16 Tanner, L. H., Lucas, S. G. & Chapman, M. G. Assessing the record and causes of
344 Late Triassic extinctions. *Earth-Science Reviews* **65**, 103-139 (2004).
- 345 17 Whiteside, J. H., Grogan, D. S., Olsen, P. E. & Kent, D. V. Climatically driven
346 biogeographic provinces of Late Triassic tropical Pangea. *Proceedings of the National*
347 *Academy of Sciences* **108**, 8972-8977, doi:10.1073/pnas.1102473108 (2011).
- 348 18 Whiteside, J., Olsen, P. E., Eglinton, T., Brookfield, M. E. & Sambrotto, R. N.
349 Compound-specific carbon isotopes from Earth's largest flood basalt eruptions
350 directly linked to the end-Triassic mass extinction. *PNAS* **107**, 6721-6725 (2010).
- 351 19 Blackburn, T. J. *et al.* Zircon U-Pb Geochronology Links the End-Triassic Extinction
352 with the Central Atlantic Magmatic Province. *Science* **340**, 941-945,
353 doi:10.1126/science.1234204 (2013).
- 354 20 Toljagić, O. & Butler, R. J. Triassic–Jurassic mass extinction as trigger for the
355 Mesozoic radiation of crocodylomorphs. *Biology Letters* **9**,
356 doi:10.1098/rsbl.2013.0095 (2013).
- 357 21 Benton, M. J. in *The Fossil Record*, vol. 2 Vol. 2 (ed M. J. Benton) 681-715
358 (Chapman and Hall, 1993).
- 359 22 Brusatte, S. L. *et al.* The origin and early radiation of dinosaurs. *Earth-Science*
360 *Reviews* **101**, 68-100 (2010).
- 361 23 Langer, M. C., Ezcurra, M. D., Bittencourt, J. S. & Novas, F. E. The origin and early
362 evolution of dinosaurs. *Biol. Rev.* **85**, 55-110, doi:10.1111/j.1469-185X.2009.00094.x
363 (2010).

- 364 24 Carrano, M. T. & Butler, R. J. Taxonomic occurrences of Triassic to Jurassic
365 Tetrapoda. Paleobiology Database (<http://paleobiodb.org>, 2013-2015).
- 366 25 Roy, K. Effects of the Mesozoic Marine Revolution on the Taxonomic, Morphologic,
367 and Biogeographic Evolution of a Group: Aporrhaid Gastropods During the
368 Mesozoic. *Paleobiology* **20**, 274-296, doi:10.2307/2401004 (1994).
- 369 26 Budd, A. F. & Coates, A. G. Nonprogressive evolution in a clade of Cretaceous
370 *Montastraea*-like corals. *Paleobiology* **18**, 425-446 (1992).
- 371 27 Jablonski, D. Larval ecology and macroevolution in marine invertebrates. *Bulletin of*
372 *Marine Science* **39**, 565-587 (1986).
- 373 28 Foote, M. Pulsed origination and extinction in the marine realm. *Paleobiology* **31**, 6-
374 20, doi:10.1666/0094-8373(2005)031<0006:poaeit>2.0.co;2 (2005).
- 375 29 Bambach, R. K. Phanerozoic biodiversity and mass extinctions. *Annual Review of*
376 *Earth and Planetary Sciences* **34**, 127-155 (2006).
- 377 30 Stubbs, T. L., Pierce, S. E., Rayfield, E. J. & Anderson, P. S. L. Morphological and
378 biomechanical disparity of crocodile-line archosaurs following the end-Triassic
379 extinction. *Proceedings of the Royal Society B: Biological Sciences* **280**,
380 doi:10.1098/rspb.2013.1940 (2013).
- 381 31 Upchurch, P., Mannion, P. D., Benson, R. B. J., Butler, R. J. & Carrano, M. T.
382 Geological and anthropogenic controls on the sampling of the terrestrial fossil record:
383 a case study from the Dinosauria. *Geological Society, London, Special Publications*
384 **358**, 209-240, doi:10.1144/sp358.14 (2011).
- 385 32 Wang, S. C. & Dodson, P. Estimating the diversity of dinosaurs. *PNAS* **103**, 13601-
386 13605 (2006).
- 387 33 Sun, Y. D. *et al.* Lethally Hot Temperatures During the Early Triassic Greenhouse.
388 *Science* **338**, 366-370, doi:10.1126/science.1224126 (2012).

- 389 34 Lloyd, G. T., Young, J. R. & Smith, A. B. Taxonomic structure of the fossil record is
390 shaped by sampling bias. *Systematic Biology* **61**, 80-89 (2012).
- 391 35 Benson, R. B. J. & Upchurch, P. Diversity trends in the establishment of terrestrial
392 vertebrate ecosystems: Interactions between spatial and temporal sampling biases.
393 *Geology* **41**, 43-46, doi:10.1130/g33543.1 (2013).
- 394 36 Dunhill, A. M., Hannisdal, B. & Benton, M. J. Disentangling geological megabias and
395 common-cause from redundancy in the British fossil record. *Nature Communications*
396 **5**, 4818, doi:10.1038/ncomms5818 (2014).
- 397 37 Benton, M. J., Dunhill, A. M., Lloyd, G. T. & Marx, F. G. Assessing the quality of the
398 fossil record: insights from vertebrates. *Geological Society, London, Special*
399 *Publications* **358**, 63-94, doi:10.1144/sp358.6 (2011).
- 400 38 Dunhill, A. M., Benton, M. J., Twitchett, R. J. & Newell, A. J. Testing the fossil
401 record: sampling proxies and scaling in the British Triassic-Jurassic.
402 *Palaeogeography Palaeoclimatology Palaeoecology* **404**, 1-11,
403 doi:http://dx.doi.org/10.1016/j.palaeo.2014.03.026 (2014).
- 404 39 Benson, R. B. J. & Butler, R. J. Uncovering the diversification history of marine
405 tetrapods: ecology influences the effect of geological sampling biases. *Geological*
406 *Society, London, Special Publications* **358**, 191-208, doi:10.1144/sp358.13 (2011).
- 407 40 Benson, R. B. J., Butler, R. J., Lindgren, J. & Smith, A. S. Mesozoic marine tetrapod
408 diversity: mass extinctions and temporal heterogeneity in geological megabiases
409 affecting vertebrates. *Proceedings of the Royal Society B* **277**, 829-834 (2010).
- 410 41 Fröbisch, J. Global taxonomic diversity of Anomodonts (Tetrapoda, Therapsida) and
411 the terrestrial rock record across the Permian-Triassic boundary. *PLoS ONE* **3**, e3733,
412 doi:10.1371/journal.pone.0003733 (2008).

413 42 Barrett, P. M., McGowan, A. J. & Page, V. Dinosaur diversity and the rock record.
414 *Proceedings of the Royal Society B* **276**, 2667-2674 (2009).

415 43 Butler, R. J., Barrett, P. M., Nowbath, S. & Upchurch, P. Estimating the effects of the
416 rock record on pterosaur diversity patterns: implications for hypotheses of
417 bird/pterosaur competitive replacement. *Paleobiology* **35**, 432-446 (2009).

418 44 Jablonski, D. Scale and hierarchy in macroevolution. *Palaeontology* **50**, 87-109,
419 doi:10.1111/j.1475-4983.2006.00615.x (2007).

420 45 Benton, M. J. The origins of modern biodiversity on land. *Phil. Trans. R. Soc. Lond.B*
421 **365**, 3667-3679 (2010).

422 46 Alroy, J. The shifting balance of diversity among major marine animal groups.
423 *Science* **329**, 1191-1194 (2010).

424 47 Sepkoski, J. J., Bambach, R. K., Raup, D. M. & Valentine, J. W. Phanerozoic marine
425 diversity and the fossil record. *Nature* **293**, 435-437 (1981).

426 48 Jablonski, D. Body-size evolution in Cretaceous molluscs and the status of Cope's
427 rule. *Nature* **385**, 250-252, doi:10.1038/385250a0 (1997).

428 49 Benson, R. B. J., Frigot, R. A., Goswami, A., Andres, B. & Butler, R. J. Competition
429 and constraint drove Cope's rule in the evolution of giant flying reptiles. *Nature*
430 *Communications* **5**, 3567, doi:10.1038/ncomms4567 (2014).

431 50 Raia, P., Carotenuto, F., Passaro, F., Fulgione, D. & Fortelius, M. Ecological
432 specialization in fossil mammals explains Cope's Rule. *American Naturalist* **179**, 328-
433 337 (2012).

434 51 Van M. Savage, James F. Gillooly, James H. Brown, Geoffrey B. West &
435 Eric L. Charnov. Effects of Body Size and Temperature on Population Growth. *The*
436 *American Naturalist* **163**, 429-441, doi:10.1086/381872 (2004).

- 437 52 Don, R. L. Influence of Body Size and Population Density on Fertilization Success
438 and Reproductive Output in a Free-Spawning Invertebrate. *Biological Bulletin* **181**,
439 261-268, doi:10.2307/1542097 (1991).
- 440 53 Cooper, N., Bielby, J., Thomas, G. H. & Purvis, A. Macroecology and extinction risk
441 correlates of frogs. *Glob. Ecol. Biogeogr.* **17**, 211-221, doi:10.1111/j.1466-
442 8238.2007.00355.x (2008).
- 443 54 Pimm, S. L. *et al.* The biodiversity of species and their rates of extinction,
444 distribution, and protection. *Science* **344**, doi:10.1126/science.1246752 (2014).
- 445 55 Cardillo, M. *et al.* The predictability of extinction: biological and external correlates
446 of decline in mammals. *Proceedings of the Royal Society B: Biological Sciences* **275**,
447 1441-1448, doi:10.1098/rspb.2008.0179 (2008).
- 448 56 Slatyer, R. A., Hirst, M. & Sexton, J. P. Niche breadth predicts geographical range
449 size: a general ecological pattern. *Ecology Letters* **16**, 1104-1114,
450 doi:10.1111/ele.12140 (2013).
- 451 57 McCain, C. M. & King, S. R. B. Body size and activity times mediate mammalian
452 responses to climate change. *Global Change Biology* **20**, 1760-1769,
453 doi:10.1111/gcb.12499 (2014).
- 454 58 Exploring macroevolution using modern and fossil data. *Proceedings of the Royal*
455 *Society B, Biological Sciences* **282**, 20150569, doi:10.1098/rspb.2015.0569 (2015).
- 456 59 Smith, A. G., Smith, D. G. & Funnell, B. M. *Atlas of Mesozoic and Cenozoic*
457 *coastlines*. 99 (Cambridge University Press, 1994).
- 458 60 Gradstein, F. M., Ogg, J. G., Schmitz, M. D. & Ogg, G. M. *The Geologic Time Scale*
459 *2012*. Vol. 2 1144 (Elsevier, 2012).
- 460 61 Haq, B. U., Hardenbol, J. & Vail, P. R. Chronology of fluctuating sea levels since the
461 Triassic. *Science* **235**, 1156-1167 (1987).

- 462 62 Butler, R. J., Benson, R. B. J., Carrano, M. T., Mannion, P. D. & Upchurch, P. Sea
463 level, dinosaur diversity and sampling biases: investigating the 'common cause'
464 hypothesis in the terrestrial realm. *Proc. R. Soc. Lond. B* **278**, 1107-1113 (2011).
- 465 63 Raup, D. M. Systematists follow the fossils. *Paleobiology* **3**, 328-329 (1977).
- 466 64 PALEOMAP Project - PointTracker v. 7h (2010).
- 467 65 Myers, C. E. & Lieberman, B. S. Sharks that pass in the night: using Geographical
468 Information Systems to investigate competition in the Cretaceous Western Interior
469 Seaway. *Proceedings of the Royal Society B: Biological Sciences* **278**, 681 (2010).
- 470 66 Myers, C. E., MacKenzie, R. A. & Lieberman, B. S. Greenhouse biogeography: the
471 relationship of geographic range to invasion and extinction in the Cretaceous Western
472 Interior Seaway. *Paleobiology* **39**, 135-148, doi:10.1666/0094-8373-39.1.135 (2013).
- 473 67 Foote, M. Origination and Extinction Components of Taxonomic Diversity: general
474 problems. *Paleobiology* **26** (supplement), 74-102, doi:10.2307/2666101 (2000).
- 475 68 Foote, M. Origination and extinction through the Phanerozoic: a new approach. *The*
476 *Journal of Geology* **111**, 125-148 (2003).
- 477 69 Foote, M. Pulsed origination and extinction in the marine realm. *Paleobiology* **31**, 6-
478 20 (2005).
- 479 70 Rode, A. L. & Lieberman, B. S. Integrating evolution and biogeography: A case study
480 involving Devonian crustaceans. *Journal of Paleontology* **79**, 267-276 (2005).

481

482 **Acknowledgements**

483 We thank Matthew Carrano, Richard Butler and various other data compilers from the
484 Paleobiology Database. This is *Paleobiology Database* official publication 232. We thank
485 Stephen Brusatte, Michael Benton, Paul Wignall, Nick Priest and Richard Butler for offering
486 valuable critical advice on earlier drafts of this manuscript. A.M.D. is supported by a Royal

Commission for the Exhibition of 1851 Fellowship. M.A.W. thanks the Leverhulme Trust for support (Grant F/00351/Z).

Author Contributions

A.M.D. designed the project and compiled the data. M.A.W. wrote scripts. A.M.D. and M.A.W. analysed the data and wrote the manuscript.

Additional Information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>.

Competing financial interests: The authors declare no competing financial interests.

Figure Legends

Figure 1| Mean rates of change in geographic range size and diversification rates for terrestrial vertebrates, partitioned by time bin. Rate of change in geographic range size (Δ Geographic range) as represented by (a) convex hulls around raw palaeogeographic occurrences and (b) mean great circle distance (GCD) between palaeogeographic occurrences standardised to 5 and 10 samples; and (c) mean vertebrate diversification rates of ranged-through diversity data (Δ Diversification). The fossil records of the Ladinian¹², Toarcian¹⁹, and much of the Middle Jurassic²⁸⁻³⁰, are of lower quality than the rest of Mesozoic, and this may contribute to some of the negative diversity changes therein. The drop in diversity observed through the Rhaetian could also be regarded as a sampling artefact as the Rhaetian is not as well sampled as the preceding Norian. However, the upper Triassic represents one of the highest quality terrestrial fossil records^{22,30}. Alternating grey and white bars correspond to Triassic-Jurassic epochs..

Figure 2| Scatter plots of diversification rates against per lineage rates of change in geographic range size. (a-c) Rate of geographic range size change (Δ Geographic range) as calculated using convex hulls around raw occurrence data for (a) all time bins; (b) Late Triassic; (c) Rhaetian. (d-f) Rate of geographic range size change calculated as mean great circle distances between occurrences standardised to samples of 5 and 10 occurrences for (d) all time bins; (e) Late Triassic; and (f) Rhaetian. Spearman rank correlation coefficients *significant at $p < 0.05$, **significant at $p < 0.01$, ***significant at $p < 0.001$..

Figure 3| Geographic range maps before and after the end-Triassic mass extinction. Convex hull geographic range maps and mean generic diversity of six terrestrial vertebrate groups during the (a) Rhaetian and (b) Hettangian. The Aetosauria, Cynognathia and Phytosauria became extinct during the biotic crisis, despite the widespread distribution of

Cynognathia and Phytosauria in the Rhaetian. The Probainognathia, Sauropodomorpha and Theropoda all survived the biotic crisis and expanded their ranges in the Hettangian, albeit with different diversification trajectories. The diversity of Sauropodomorpha increased, the diversity of Theropoda was stationary, and that of Probainognathia decreased across the boundary.

Figure 4| Sampling and environmental proxy data. (a) non-marine area⁵⁸; (b) average sea level⁶⁰; (c) terrestrial formation count²⁴; (d) fossil abundance²⁴; (e) total geographic range of all taxa. Alternating grey and white bars correspond to Triassic-Jurassic epochs.

Table 1| Summary of multiple regression models of diversity change (dependent variable) in terms of geographic range change and sampling and environmental proxies.

Model	Dependent	Independents	adj R ²	p	AIC
Convex full	Diversity change	Range change + land area + sea level + formations + abundance + total range	0.53	< 0.001	-98.59
Convex best	Diversity change	Range change	0.64	< 0.001	-100.53
GCD5 full	Diversity change	Range change + land area + sea level + formations + abundance + total range	0.22	< 0.001	-65.5
GCD5 best	Diversity change	Range change + total range + sea level	0.23	< 0.001	-68.95
GCD10 full	Diversity change	Range change + land area + sea level + formations + abundance + total range	0.32	< 0.001	-63.66
GCD10 best	Diversity change	Range change + total range + sea level	0.35	< 0.001	-68.97

538







